


Chapter 20: ENTROPY AND THE SECOND LAW OF THERMODYNAMICS

- In a reversible process the system:
 - is always close to equilibrium states
 - is close to equilibrium states only at the beginning and end
 - might never be close to any equilibrium state
 - is close to equilibrium states throughout, except at the beginning and end
 - is none of the aboveans: A
- A slow (quasi-static) process is NOT reversible if:
 - the temperature changes
 - energy is absorbed or emitted as heat
 - work is done on the system
 - friction is present
 - the pressure changesans: D
- The difference in entropy $\Delta S = S_B - S_A$ for two states A and B of a system can be computed as the integral $\int dQ/T$ provided:
 - A and B are on the same adiabat
 - A and B have the same temperature
 - a reversible path is used for the integral
 - the change in internal energy is first computed
 - the energy absorbed as heat by the system is first computedans: C
- Possible units of entropy are:
 - J
 - J/K
 - J⁻¹
 - liter·atm
 - cal/molans: B
- Which of the following is NOT a state variable?
 - Work
 - Internal energy
 - Entropy
 - Temperature
 - Pressureans: A

6. The change in entropy is zero for:
- A. reversible adiabatic pr
 - B. reversible isothermal processes
 - C. reversible processes during which no work is done
 - D. reversible isobaric processes
 - E. all adiabatic processes
- ans: A
7. Which of the following processes leads to a change in entropy of zero for the system undergoing the process?
- A. Non-cyclic isobaric (constant pressure)
 - B. Non-cyclic isochoric (constant volume)
 - C. Non-cyclic isothermal (constant temperature)
 - D. Any closed cycle
 - E. None of these
- ans: D
8. Rank, from smallest to largest, the changes in entropy of a pan of water on a hot plate, as the temperature of the water
- 1. goes from 20° C to 30° C
 - 2. goes from 30° C to 40° C
 - 3. goes from 40° C to 45° C
 - 4. goes from 80° C to 85° C
- A. 1, 2, 3, 4
 - B. 4, 3, 2, 1
 - C. 1 and 2 tie, then 3 and 4 tie
 - D. 3 and 4 tie, then 1 and 2 tie
 - E. 4, 3, 2, 1
- ans: E
9. An ideal gas expands into a vacuum in a rigid vessel. As a result there is:
- A. a change in entropy
 - D. an increase of pressure
 - B. a change in temperature
 - E. a decrease of internal energy
 - C. a change in phase
- ans: A
10. Consider all possible isothermal contractions of an ideal gas. The change in entropy of the gas:
- A. is zero for all of them
 - B. does not decrease for any of them
 - C. does not increase for any of them
 - D. increases for all of them
 - E. decreases for all of them
- ans: E

11. An ideal gas is taken reversibly from state i , at temperature T_1 , to any of the other states labeled I, II, III, IV, and V. Rank the five processes according to the change in entropy of the gas, least to greatest.



- A. I, II, III, IV, V
 B. V, IV, III, II, I
 C. I, then II, III, IV, and V tied
 D. I, II, III, and IV tied, then V
 E. I and V tied, then II, III, IV
 ans: A
12. An ideal gas, consisting of n moles, undergoes a reversible isothermal process during which the volume changes from V_i to V_f . The change in entropy of the thermal reservoir in contact with the gas is given by:
- A. $nR(V_f - V_i)$
 B. $nR \ln(V_f - V_i)$
 C. $nR \ln(V_i/V_f)$
 D. $nR \ln(V_f/V_i)$
 E. none of the above (entropy can't be calculated for a reversible process)
 ans: C
13. One mole of an ideal gas expands reversibly and isothermally at temperature T until its volume is doubled. The change of entropy of this gas for this process is:
- A. $R \ln 2$
 B. $(\ln 2)/T$
 C. 0
 D. $RT \ln 2$
 E. $2R$
 ans: A

14. An ideal gas, consisting of n moles, undergoes an irreversible process in which the temperature has the same value at the beginning and the end. The change in entropy of the gas is given by.
- $nR(V_f - V_i)$
 - $nR \ln(V_f - V_i)$
 - $nR \ln(V_i/V_f)$
 - $nR \ln(V_f/V_i)$
 - none of the above (entropy can't be calculated for an irreversible process)
- ans: D
15. The temperature of n moles of a gas is increased from T_i to T_f at constant volume. If the molar specific heat at constant volume is C_V and is independent of temperature, then change in the entropy of the gas is:
- $nC_V \ln(T_f/T_i)$
 - $nC_V \ln(T_i/T_f)$
 - $nC_V \ln(T_f - T_i)$
 - $nC_V \ln(1 - T_i/T_f)$
 - $nC_V(T_f - T_i)$
- ans: A
16. Consider the following processes: The temperature of two identical gases are increased from the same initial temperature to the same final temperature. Reversible processes are used. For gas A the process is carried out at constant volume while for gas B it is carried out at constant pressure. The change in entropy:
- is the same for A and B
 - is greater for A
 - is greater for B
 - is greater for A only if the initial temperature is low
 - is greater for A only if the initial temperature is high
- ans: C
17. A hot object and a cold object are placed in thermal contact and the combination is isolated. They transfer energy until they reach a common temperature. The change ΔS_h in the entropy of the hot object, the change ΔS_c in the entropy of the cold object, and the change ΔS_{total} in the entropy of the combination are:
- $\Delta S_h > 0, \Delta S_c > 0, \Delta S_{\text{total}} > 0$
 - $\Delta S_h < 0, \Delta S_c > 0, \Delta S_{\text{total}} > 0$
 - $\Delta S_h < 0, \Delta S_c > 0, \Delta S_{\text{total}} < 0$
 - $\Delta S_h > 0, \Delta S_c < 0, \Delta S_{\text{total}} > 0$
 - $\Delta S_h > 0, \Delta S_c < 0, \Delta S_{\text{total}} < 0$
- ans: B

18. Let S_I denote the change in entropy of a sample for an irreversible process from state A to state B. Let S_R denote the change in entropy of a sample for a reversible process from state A to state B. Then:

- A. $S_I > S_R$
- B. $S_I = S_R$
- C. $S_I < S_R$
- D. $S_I = 0$
- E. $S_R = 0$

ans: B

19. For all adiabatic processes:

- A. the entropy of the system does not change
- B. the entropy of the system increases
- C. the entropy of the system decreases
- D. the entropy of the system does not increase
- E. the entropy of the system does not decrease

ans: E

20. For all reversible processes involving a system and its environment:

- A. the entropy of the system does not change
- B. the entropy of the system increases
- C. the total entropy of the system and its environment does not change
- D. the total entropy of the system and its environment increases
- E. none of the above

ans: C

21. For all irreversible processes involving a system and its environment:

- A. the entropy of the system does not change
- B. the entropy of the system increases
- C. the total entropy of the system and its environment does not change
- D. the total entropy of the system and its environment increases
- E. none of the above

ans: D

22. According to the second law of thermodynamics:

- A. heat energy cannot be completely converted to work
- B. work cannot be completely converted to heat energy
- C. for all cyclic processes we have $dQ/T < 0$
- D. the reason all heat engine efficiencies are less than 100% is friction, which is unavoidable
- E. all of the above are true

ans: A

23. Consider the following processes:
- I. Energy flows as heat f
 - II. Work is done on a system and an equivalent amount of energy is rejected as heat by the system
 - III. Energy is absorbed as heat by a system and an equivalent amount of work is done by the system
- Which are never found to occur?
- A. Only I
 - B. Only II
 - C. Only III
 - D. Only II and III
 - E. I, II, and III
- ans: C
24. An inventor suggests that a house might be heated by using a refrigerator to draw energy as heat from the ground and reject energy as heat into the house. He claims that the energy supplied to the house as heat can exceed the work required to run the refrigerator. This:
- A. is impossible by first law
 - B. is impossible by second law
 - C. would only work if the ground and the house were at the same temperature
 - D. is impossible since heat energy flows from the (hot) house to the (cold) ground
 - E. is possible
- ans: E
25. In a thermally insulated kitchen, an ordinary refrigerator is turned on and its door is left open. The temperature of the room:
- A. remains constant according to the first law of thermodynamics
 - B. increases according to the first law of thermodynamics
 - C. decreases according to the first law of thermodynamics
 - D. remains constant according to the second law of thermodynamics
 - E. increases according to the second law of thermodynamics
- ans: B
26. A heat engine:
- A. converts heat input to an equivalent amount of work
 - B. converts work to an equivalent amount of heat
 - C. takes heat in, does work, and loses energy as heat
 - D. uses positive work done on the system to transfer heat from a low temperature reservoir to a high temperature reservoir
 - E. uses positive work done on the system to transfer heat from a high temperature reservoir to a low temperature reservoir.
- ans: C

27. A heat engine absorbs energy of magnitude $|Q_H|$ as heat from a high temperature reservoir, does work of magnitude $|W|$, and rejects energy of magnitude $|Q_L|$ to a low temperature reservoir. Its efficiency is:

- A. $|Q_H|/|W|$
- B. $|Q_L|/|W|$
- C. $|Q_H|/|Q_L|$
- D. $|W|/|Q_H|$
- E. $|W|/|Q_L|$

ans: D

28. The temperatures T_C of the cold reservoirs and the temperatures T_H of the hot reservoirs for four Carnot heat engines are

engine 1: $T_C = 400$ K and $T_H = 500$ K

engine 2: $T_C = 500$ K and $T_H = 600$ K

engine 3: $T_C = 400$ K and $T_H = 600$ K

engine 4: $T_C = 600$ K and $T_H = 800$ K

Rank these engines according to their efficiencies, least to greatest

- A. 1, 2, 3, 4
- B. 1 and 2 tie, then 3 and 4 tie
- C. 2, 1, 3, 4
- D. 1, 2, 4, 3
- E. 2, 1, 4, 3

ans: E

29. A Carnot heat engine runs between a cold reservoir at temperature T_C and a hot reservoir at temperature T_H . You want to increase its efficiency. Of the following, which change results in the greatest increase in efficiency? The value of ΔT is the same for all changes.

- A. Raise the temperature of the hot reservoir by ΔT
- B. Raise the temperature of the cold reservoir by ΔT
- C. Lower the temperature of the hot reservoir by ΔT
- D. Lower the temperature of the cold reservoir by ΔT
- E. Lower the temperature of the hot reservoir by $\frac{1}{2}\Delta T$ and raise the temperature of the cold reservoir by $\frac{1}{2}\Delta T$

ans: D

30. 31. A certain heat engine draws 500 cal/s from a water bath at 27° C and transfers 400 cal/s to a reservoir at a lower temperature. The efficiency of this engine is:

- A. 80%
- B. 75%
- C. 55%
- D. 25%
- E. 20%

ans: E

32. A heat engine that in each cycle does positive work and loses energy as heat, with no heat energy input, would violate
- A. the zeroth law of thermodynamics
 - B. the first law of thermodynamics
 - C. the second law of thermodynamics
 - D. the third law of thermodynamics
 - E. Newton's second law
- ans: B
33. A cyclical process that transfers energy as heat from a high temperature reservoir to a low temperature reservoir with no other change would violate:
- A. the zeroth law of thermodynamics
 - B. the first law of thermodynamics
 - C. the second law of thermodynamics
 - D. the third law of thermodynamics
 - E. none of the above
- ans: E
34. On a warm day a pool of water transfers energy to the air as heat and freezes. This is a direct violation of:
- A. the zeroth law of thermodynamics
 - B. the first law of thermodynamics
 - C. the second law of thermodynamics
 - D. the third law of thermodynamics
 - E. none of the above
- ans: C
35. A heat engine in each cycle absorbs energy of magnitude $|Q_H|$ as heat from a high temperature reservoir, does work of magnitude $|W|$, and then absorbs energy of magnitude $|Q_L|$ as heat from a low temperature reservoir. If $|W| = |Q_H| + |Q_L|$ this engine violates:
- A. the zeroth law of thermodynamics
 - B. the first law of thermodynamics
 - C. the second law of thermodynamics
 - D. the third law of thermodynamics
 - E. none of the above
- ans: C
36. A heat engine in each cycle absorbs energy from a reservoir as heat and does an equivalent amount of work, with no other changes. This engine violates:
- A. the zeroth law of thermodynamics
 - B. the first law of thermodynamics
 - C. the second law of thermodynamics
 - D. the third law of thermodynamics
 - E. none of the above
- ans: C

37. A Carnot cycle:
- A. is bounded by two isot
 - B. consists of two isothermal and two constant volume processes
 - C. is any four-sided process on a p - V graph
 - D. only exists for an ideal gas
 - E. has an efficiency equal to the enclosed area on a p - V diagram
- ans: A
38. According to the second law of thermodynamics:
- A. all heat engines have the same efficiency
 - B. all reversible heat engines have the same efficiency
 - C. the efficiency of any heat engine is independent of its working substance
 - D. the efficiency of a Carnot engine depends only on the temperatures of the two reservoirs
 - E. all Carnot engines theoretically have 100% efficiency
- ans: D
39. A Carnot heat engine operates between 400 K and 500 K. Its efficiency is:
- A. 20%
 - B. 25%
 - C. 44%
 - D. 79%
 - E. 100%
- ans: A
40. A Carnot heat engine operates between a hot reservoir at absolute temperature T_H and a cold reservoir at absolute temperature T_C . Its efficiency is:
- A. T_H/T_C
 - B. T_C/T_H
 - C. $1 - T_H/T_C$
 - D. $1 - T_C/T_H$
 - E. 100%
- ans: D
41. A heat engine operates between a high temperature reservoir at T_H and a low temperature reservoir at T_L . Its efficiency is given by $1 - T_L/T_H$:
- A. only if the working substance is an ideal gas
 - B. only if the engine is reversible
 - C. only if the engine is quasi-static
 - D. only if the engine operates on a Stirling cycle
 - E. no matter what characteristics the engine has
- ans: B

42. The maximum theoretical efficiency of a Carnot heat engine operating between reservoirs at the steam point and at room temperature is
- A. 10%
 - B. 20%
 - C. 50%
 - D. 80%
 - E. 99%

ans: B

43. An inventor claims to have a heat engine that has an efficiency of 40% when it operates between a high temperature reservoir of 150°C and a low temperature reservoir of 30°C . This engine:
- A. must violate the zeroth law of thermodynamics
 - B. must violate the first law of thermodynamics
 - C. must violate the second law of thermodynamics
 - D. must violate the third law of thermodynamics
 - E. does not necessarily violate any of the laws of thermodynamics

ans: C

44. A Carnot heat engine and an irreversible heat engine both operate between the same high temperature and low temperature reservoirs. They absorb the same energy from the high temperature reservoir as heat. The irreversible engine:
- A. does more work
 - B. transfers more energy to the low temperature reservoir as heat
 - C. has the greater efficiency
 - D. has the same efficiency as the reversible engine
 - E. cannot absorb the same energy from the high temperature reservoir as heat without violating the second law of thermodynamics

ans: B

45. A perfectly reversible heat pump with a coefficient of performance of 14 supplies energy to a building as heat to maintain its temperature at 27°C . If the pump motor does work at the rate of 1 kW, at what rate does the pump supply energy to the building as heat?
- A. 15 kW
 - B. 3.85 kW
 - C. 1.35 kW
 - D. 1.07 kW
 - E. 1.02 kW

ans: A

46. A heat engine operates between 200 K and 100 K. In each cycle it takes 100 J from the hot reservoir, loses 25 J to the cold reservoir, and does 75 J of work. This heat engine violates:
- A. both the first and second laws of thermodynamics
 - B. the first law but not the second law of thermodynamics
 - C. the second law but not the first law of thermodynamics
 - D. neither the first law nor the second law of thermodynamics
 - E. cannot answer without knowing the mechanical equivalent of heat

ans: C

47. A refrigerator absorbs energy of magnitude $|Q_C|$ as heat from a low temperature reservoir and transfers energy of magnitude $|Q_H|$ to a high temperature reservoir. The coefficient of performance is given by:
- $|Q_C|/W$
 - $|Q_H|/W$
 - $(|Q_C| + |Q_H|)/W$
 - $W/|Q_C|$
 - $W/|Q_H|$
- ans: A
48. A reversible refrigerator operates between a low temperature reservoir at T_C and a high temperature reservoir at T_H . Its coefficient of performance is given by:
- $(T_H - T_C)/T_C$
 - $T_C/(T_H - T_C)$
 - $(T_H - T_C)/T_H$
 - $T_H/(T_H - T_C)$
 - $T_H(T_H + T_C)$
- ans: B
49. An Carnot refrigerator runs between a cold reservoir at temperature T_C and a hot reservoir at temperature T_H . You want to increase its coefficient of performance. Of the following, which change results in the greatest increase in the coefficient? The value of ΔT is the same for all changes.
- Raise the temperature of the hot reservoir by ΔT
 - Raise the temperature of the cold reservoir by ΔT
 - Lower the temperature of the hot reservoir by ΔT
 - Lower the temperature of the cold reservoir by ΔT
 - Lower the temperature of the hot reservoir by $\frac{1}{2}\Delta T$ and raise the temperature of the cold reservoir by $\frac{1}{2}\Delta T$
- ans: B
50. For one complete cycle of a reversible heat engine, which of the following quantities is NOT zero?
- the change in the entropy of the working gas
 - the change in the pressure of the working gas
 - the change in the internal energy of the working gas
 - the work done by the working gas
 - the change in the temperature of the working gas
- ans: D

51. Twenty-five identical molecules are in a box. Microstates are designated by identifying the molecules in the left and right halves of the box. The Boltzmann constant is 1.38×10^{-23} J/K. The entropy associated with the configuration for which 15 molecules are in the left half and 10 molecules are in the right half is:

A. 1.03×10^{23}
B. 3.27×10^6
C. 150
D. 25
E. 5

ans: B

52. Twenty-five identical molecules are in a box. Microstates are designated by identifying the molecules in the left and right halves of the box. The Boltzmann constant is 1.38×10^{-23} J/K. The entropy associated with the configuration for which 15 molecules are in the left half and 10 molecules are in the right half is:

A. 2.07×10^{-22} J/K
B. 7.31×10^{-22} J/K
C. 4.44×10^{-23} J/K
D. 6.91×10^{-23} J/K
E. 2.22×10^{-23} J/K

ans: A

53. The thermodynamic state of a gas changes from one with 3.8×10^{18} microstates to one with 7.9×10^{19} microstates. The Boltzmann constant is 1.38×10^{-23} J/K. The change in entropy is:

A. $\Delta S = 0$
B. $\Delta S = 1.04 \times 10^{-23}$ J/K
C. $\Delta S = -1.04 \times 10^{-23}$ J/K
D. $\Delta S = 4.19 \times 10^{-23}$ J/K
E. $\Delta S = -4.19 \times 10^{-23}$ J/K

ans: D

54. Let k be the Boltzmann constant. If the configuration of the molecules in a gas changes so that the multiplicity is reduced to one-third its previous value, the entropy of the gas changes by:

A. $\Delta S = 0$
B. $\Delta S = 3k \ln 2$
C. $\Delta S = -3k \ln 2$
D. $\Delta S = -k \ln 3$
E. $\Delta S = k \ln 3$

ans: D

55. Let k be the Boltzmann constant. If the configuration of molecules in a gas changes from one with a multiplicity of M_1 to one with a multiplicity of M_2 , then entropy changes by:

A. $\Delta S = 0$
B. $\Delta S = k(M_2 - M_1)$
C. $\Delta S = kM_2/M_1$
D. $\Delta S = k \ln(M_2 M_1)$
E. $\Delta S = k \ln(M_2/M_1)$

ans: E

56. Let k be the Boltzmann constant. If the thermodynamic state of a gas at temperature T changes isothermally and its volume is tripled, the energy input to the gas as heat is:
- A. $Q = 0$
 - B. $Q = 3kT$
 - C. $Q = -3kT$
 - D. $kT \ln 3$
 - E. $-kT \ln 3$
- ans: D

